

Craig Milton
Policy Manager
NSW Minerals Council

Via Email: cmilton@nswmining.com.au

26 April 2024

Upper Hunter Air Quality Monitoring Network Analysis Project – Annual Review for 2023

Dear Craig,

1 Introduction

Zephyr Environmental Pty Ltd (Zephyr) has been commissioned by the NSW Minerals Council (NSWMC) Upper Hunter Mining Dialogue (UHMD) to provide an updated air quality analysis, now incorporating the calendar year (CY) 2023, for the Upper Hunter following continued interest in air quality in the region.

The analysis of CY2023 air quality data follows on from the previous work prepared on air quality trends across NSW namely:

- Air Quality Monitoring Data Analysis Project ('the AQ Data Analysis Project', dated 16 November 2020).
- Upper Hunter Air Quality Monitoring Network Analysis Project ('0012 UHMD UHAQMN Annual Data Review L1 Final Update', dated 9 December 2021)
- Upper Hunter Air Quality Monitoring Network Analysis Project – Annual Review for 2021 and 2022 ('0130 NSWMC UHAQMP Annual Data Review Final, dated 26 October 2023)

Section 1 of the AQ Data Analysis Project provides background and context to this work, and confirms that the original project was designed to answer two specific air quality questions:

1. Has the air quality in the Upper Hunter Valley changed since monitoring began? and
2. Is the air quality in the Upper Hunter Valley measured at the monitoring stations different from air quality measured at other locations in NSW?

Section 2 of the AQ Data Analysis Project provides an overview of particulate matter size fractions and ambient air quality criteria, including definitions of the terms PM₁₀ and PM_{2.5} (particulate matter less than 10 and 2.5 micrometres in aerodynamic diameter, respectively).

Section 3 of the AQ Data Analysis Project provides detail on the 14 air quality monitoring stations that comprise the Upper Hunter Air Quality Monitoring Network (UHAQMN) and their grouping to reflect the following purposes:

- Larger populations (Muswellbrook, Singleton and Aberdeen)
- Smaller communities (Bulga, Camberwell, Jerrys Plains, Maison Dieu, Warkworth, Wybong)
- Diagnostic (Mount Thorley, Muswellbrook NW, Singleton NW)
- Background (Merriwa, Singleton South)

Section 4 and Section 5 of the AQ Data Analysis Project provide a summary of the air quality monitoring data collected to date and their analysis respectively.

The focus of this letter is to provide an update to the AQ Data Analysis Project, now including data from CY2023.

The conclusions from last years' review of CY2021 and CY2022 data were as follows:

- As noted within the previous reports and reinforced with the inclusion of CY2021 and CY2022 data, the temporal trends in the UHAQMN and 'Remainder of NSW' monitoring subsets show that the changes in PM₁₀ concentrations within the Upper Hunter are generally consistent with changes experienced across the rest of NSW.
- With the inclusion of CY2021 and CY2022 data, it continues to be seen that there is a negative correlation between rainfall and particulate matter concentrations across the UHAQMN. Given the consistency between PM₁₀ trends across NSW and the UHAQMN, this relationship also holds for NSW PM₁₀ concentrations more broadly.
- The differences between PM₁₀ concentrations at Background stations and Diagnostic stations are near identical across CY2013-CY2022 (up to 2 µg/m³ variability), while the range in annual average PM₁₀ concentrations across this period is 22 µg/m³. This reinforces that changes in Upper Hunter PM₁₀ concentrations are associated with regional conditions and are indicative of a minimal change in the contribution from local emission sources inclusive of mining.
- There is no visually apparent correlation between raw coal production and ambient PM₁₀ concentrations measured by the UHAQMN between CY2013 and CY2019. The supporting figure shows inconsistency in the scale of variability in each metric across this period, with minor proportional variability in coal production relative to significant proportional variability in annual average PM₁₀.
- From 2019 to 2022, both coal production and annual average PM₁₀ concentrations have reduced year on year. However, the reductions in annual average PM₁₀ concentrations are anticipated to be related to the lower ambient temperatures and significant amounts of rainfall during these years, rather than the reduced coal production. It is recommended that this apparent trend be reevaluated in future years.

2 Monitoring Data Update

The NSW Department of Climate Change, Energy, the Environment and Water (DCCEEW), formerly NSW Department of Planning and Environment (DPE), makes available ambient air quality monitoring data for NSW via their data download facility. Data has been gathered for the most recent complete calendar year of CY2023.

Table 2.1 and Table 2.2 present a summary of annual and period average PM₁₀ and PM_{2.5} monitoring results (respectively) that have been averaged into regions/groups.

Results have been shaded using a green to red colour relative gradient scheme with lowest values shown in green, and highest values shown in red, with the median value shown in yellow. This gradient scheme has been applied to the annual data and 'all years' groups separately.

For the average across all years, the highest concentrations of PM₁₀ are measured at the UHAQMN Diagnostic stations, with the next highest observed at the Lower Hunter & Central Coast stations. The highest concentrations of PM_{2.5} are measured at the UHAQMN Large Population stations, with the next highest observed at the Sydney north-west stations. The lowest concentrations are measured at the Central Tablelands for PM₁₀ and Illawarra for PM_{2.5}, respectively.

Annual PM₁₀ and PM_{2.5} averages have increased from CY2022 to CY2023, across all regions / groups. The CY2023 annual average PM₁₀ and PM_{2.5} concentrations are generally the highest observed

concentrations since 2020. The CY2023 averages for PM₁₀ remain below the 'all years' average for all regions/groups with the exceptions of Sydney north-west (CY2023 = 17.8 µg/m³ and 'All Years' = 17.3 µg/m³) and UHAQMN – SC (CY2023 = 21.3 µg/m³ and 'All Years' = 20.9 µg/m³). The CY2023 averages for PM_{2.5} remain below the 'all years' average for all regions/groups.

Table 2.1: Annual and period average PM₁₀ concentrations by region/group and year (µg/m³)

Region / Group	Year											All years
	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	
Central tablelands	15.1	14.6	13.4	13.3	14.1	18.8	27.4	17.0	11.4	8.7	12.5	15.1
Illawarra	16.9	17.1	16.2	17.4	18.0	20.1	22.5	19.1	15.3	13.2	16.9	17.5
Lower Hunter & Central Coast	20.2	18.2	21.7	22.0	22.9	25.2	29.1	22.3	19.2	17.6	20.9	21.8
North-west slopes	16.6	15.8	14.1	15.3	15.3	20.1	33.7	16.8	12.7	10.6	15.1	16.9
South-west slopes	10.0	18.3	17.3	17.9	18.2	23.6	29.4	21.7	16.0	12.4	15.5	18.2
Sydney east	17.9	17.3	16.8	17.2	18.3	20.2	23.6	19.2	15.9	13.5	16.1	17.8
Sydney north-west	17.5	16.6	15.1	17.0	17.0	20.3	24.9	18.7	15.7	11.9	17.8	17.5
Sydney south-west	16.3	16.0	14.8	15.6	16.1	18.9	23.3	17.2	13.8	11.1	15.2	16.2
UHAQMN - BG	17.6	16.8	15.1	15.8	16.8	21.1	29.3	19.0	14.1	12.6	16.8	17.7
UHAQMN - DG	23.2	21.1	19.1	20.4	22.2	29.0	34.9	21.7	16.7	14.6	21.9	22.3
UHAQMN - LP	21.1	20.1	17.9	18.0	20.0	24.5	31.3	20.3	16.2	14.5	18.8	20.2
UHAQMN - SC	21.4	20.1	17.7	18.6	20.7	25.4	33.4	21.2	16.4	14.2	21.3	20.9

Note: UHAQMN – upper hunter air quality monitoring network, BG - background, DG – diagnostic, LP – larger populations, SC – smaller communities

Colour Coding by Percentile

0% (min.)	10%	20%	30%	40%	50% (median)	60%	70%	80%	90%	100% (max.)
-----------	-----	-----	-----	-----	--------------	-----	-----	-----	-----	-------------

Note: colour coding is applied to annual data by region (horizontally), whereas 'All years' colour coding is applied vertically, to allow comparison of data between regions.

Table 2.2: Annual and period average PM_{2.5} concentrations by region / group and year (µg/m³)

Region / Group	Year											All years
	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	
Illawarra	7.7	7.0	7.0	7.3	6.9	7.1	11.1	7.2	5.3	4.3	5.3	6.9
Lower Hunter & Central Coast	7.5	7.0	7.5	7.8	7.7	8.2	17.3	7.6	6.3	5.5	6.8	8.1
South-west slopes	7.9	7.5	7.6	7.4	8.1	8.4	11.3	10.9	6.3	5.3	6.6	7.9
Sydney east	8.2	8.4	8.3	8.1	8.4	8.2	16.5	8.0	6.9	5.2	7.1	8.5
Sydney north-west	8.3	6.7	8.0	8.3	7.4	8.3	20.5	8.2	6.9	5.1	7.1	8.6
Sydney south-west	8.0	7.5	7.4	7.6	7.8	8.7	18.9	7.9	7.0	5.0	6.8	8.4
UHAQMN - LP	8.7	8.8	8.2	8.2	8.8	8.8	18.0	8.9	6.8	5.7	7.1	8.9
UHAQMN - SC	8.2	7.8	7.2	7.5	7.4	8.4	17.3	7.5	5.7	4.8	6.1	7.4

Note: UHAQMN – upper hunter air quality monitoring network, LP – larger populations, SC – smaller communities

Colour Coding by Percentile

0% (min.)	10%	20%	30%	40%	50% (median)	60%	70%	80%	90%	100% (max.)
-----------	-----	-----	-----	-----	--------------	-----	-----	-----	-----	-------------

Note: colour coding is applied to annual data by region (horizontally), whereas 'All years' colour coding is applied vertically, to allow comparison of data between regions.

3 Analysis Update

Provided below are revisions of some of the key data tables and figures from the AQ Data Analysis Project, updated to include data from CY2023. Additional commentary is provided as it relates to the inclusion of the CY2023 data.

Comparison of UHAQMN and remainder of NSW

Table 3.1 presents a comparison of average PM₁₀ concentrations measured across NSW, with the UHAQMN and the remainder of NSW shown separately. Annual data have also been presented as a percentage of the respective 2013-2023 average.

This relationship has also been shown for PM_{2.5}, which has a lesser association with mechanically generated particulate emissions such as those from mining. Figure 3.1 provides a graphical representation of these data.

As noted within the AQ Data Analysis Project and reinforced with the inclusion of CY2021, CY2022 and CY2023 data, the consistency of temporal trends in the UHAQMN and 'Remainder of NSW' monitoring subsets show that the changes in PM₁₀ concentrations within the Upper Hunter are generally consistent with changes experienced across the rest of NSW.



This in turn indicates that the changes in annual average PM₁₀ concentrations are associated with regional particulate sources and that the contribution of mining operations on the UHAQMN to these trends is not discernible.

Notably, the CY2023 PM₁₀ data for the UHAQMN is below the 'all years' average. The recorded concentration is higher than values recorded in the particularly high rainfall years of CY2021 and CY2022 but remains substantially lower than the anomalously high CY2019 value associated with the 'Black Summer' bushfire event and even CY2020. A similar pattern is also observed in PM_{2.5}, indicating that changes in PM_{2.5} (less likely to be attributed to mining operations) are generally consistent with those observed elsewhere in NSW. As with the PM₁₀ data set, the moderate values for CY2023 emphasise that the annual average for CY2019 was anomalously high.

Table 3.1: Comparison of PM₁₀ and PM_{2.5} variability – UHAQMN vs remainder of NSW regions

Monitoring subset	Parameter	Year											All Years
		2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	
PM₁₀													
UHAQMN*	Concentration (µg/m ³)	21.2	20.1	17.8	18.3	20.4	25.0	32.7	20.9	16.3	14.3	20.4	20.7
	% of average (all years)	102%	97%	86%	88%	99%	121%	158%	101%	79%	69%	99%	-
Remainder of NSW	Concentration (µg/m ³)	17.4	16.7	16.2	17.0	17.5	20.9	26.0	19.5	15.7	13.3	17.2	18.0
	% of average (all years)	97%	93%	90%	94%	97%	116%	144%	108%	87%	74%	96%	-
PM_{2.5}													
UHAQMN*	Concentration (µg/m ³)	7.9	7.3	7.6	7.7	7.7	8.1	11.2	8.4	6.4	5.4	6.8	7.7
	% of average (all years)	102%	94%	98%	99%	99%	104%	144%	108%	83%	69%	88%	-
Remainder of NSW	Concentration (µg/m ³)	7.9	7.4	7.6	7.8	7.7	8.2	11.2	8.1	6.4	5.1	6.6	7.6
	% of average (all years)	102%	96%	98%	101%	99%	106%	145%	104%	82%	66%	87%	-

Note: UHAQMN – upper hunter air quality monitoring network, * - Larger Populations and Smaller Communities station groups.



Note: UHAQMN data relates to Larger Populations and Smaller Communities station groups.

Figure 3.1: Comparison of PM₁₀ (top) and PM_{2.5} (bottom) variability – UHAQMN vs remainder of NSW regions

Comparison across sub-divisions of the UHAQMN

For each station group, Table 3.2 shows the annual PM₁₀ variance against the CY2013 – CY2023 average PM₁₀ concentration for that station group. This is instructive in showing changes in the difference between station groups across the study period, into CY2023. The same data is shown graphically in Figure 3.2.

Table 3.2: Annual variance against CY2013 – CY2023 station group average (µg/m³)

Station Group	Year										
	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023
Larger Population	1	0	-2	-2	0	5	11	0	-4	-6	-1
Smaller Communities	0	-1	-3	-2	0	4	12	0	-5	-7	0
Diagnostic	1	-1	-3	-2	0	5	13	0	-5	-7	0
Background	0	-1	-3	-2	-1	3	11	1	-4	-5	-1




Figure 3.2: Comparison of trends between each UHAQMN station group

Figure 3.2 shows that the differences between PM₁₀ concentrations at Background stations and Diagnostic stations are near identical across CY2013-CY2023 (up to 1 µg/m³ variability), while the range in annual average PM₁₀ concentrations across this period is 22 µg/m³. This reinforces that changes in Upper Hunter PM₁₀ concentrations are associated with regional conditions and are indicative of a minimal change in the contribution from local emission sources inclusive of mining.

Comparison of UHAQMN and Bulga, Mount Thorley, Warkworth stations

The annual average PM₁₀ concentrations measured across the Bulga, Mount Thorley and Warkworth monitoring stations have been compared against the UHAQMN for 2018 to 2023. These three stations and the UHAQMN show concentrations increasing from 2018 to 2019 associated with the ‘Black Summer’ bushfire events. All three sites show a reduction in concentrations year on year from 2019 through to 2022, due to higher-than-average rainfall volumes particularly in CY21-CY22, and then an increase in concentrations from 2022 to 2023. The UHAQMN follows the same trend.

Comparison of UHAQMN annual average PM₁₀ and NSW/ACT rainfall

In Appendix A1.4 of the AQ Data Analysis Project, the conclusion is that there is a statistically significant relationship between NSW mean annual rainfall and annual average PM₁₀. This was further supported by the Annual Review for 2021 and 2022, which continued to show a negative correlation between rainfall and PM₁₀ across the UHAQMN.


Table 3.3 presents NSW/ACT annual rainfall and UHAQMN annual average PM₁₀ over the period CY2013-CY2023. These data are shown in Figure 3.3.

Table 3.3: NSW/ACT annual rainfall and UHAQMN annual average PM₁₀

Parameter	Year										
	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023
UHAQMN PM ₁₀ (µg/m ³)	21	20	18	18	20	25	33	21	16	14	20
NSW Rainfall (mm)	464	467	541	661	453	333	250	639	721	864	429

Figure 3.3 clearly shows that there is a negative correlation between rainfall and particulate matter concentrations across the UHAQMN. Given the consistency between PM₁₀ trends across NSW and the UHAQMN (refer Figure 3.1), this relationship also holds for NSW PM₁₀ concentrations more broadly.

CY2023 data reinforces this negative correlation. PM₁₀ shows a moderate increase in line with a moderate decrease in rainfall, compared to very high rainfall conditions and corresponding low PM₁₀ during CY2021-CY2022 (La Niña conditions).



**Figure 3.3: NSW/ACT annual rainfall and UHAQMN annual average PM₁₀
Comparison of UHAQMN annual average PM₁₀ and Hunter Valley Coal Production**

Appendix A1.1 of the AQ Data Analysis Project demonstrated that, over the period 2013-2019, the relationship between annual coal production and annual average PM₁₀ is not statistically significant. Similarly, Appendix A1.2 of the AQ Data Analysis Project showed that the relationship between annual National Pollutant Inventory (NPI) reported PM₁₀ emissions and annual average PM₁₀ concentrations is not statistically significant. The CY2021-CY2022 Annual Review found that both coal production and annual average PM₁₀ concentrations had decreased with each year from 2019 to 2022. It was determined that the reduction in annual average PM₁₀ concentrations was more likely to be related to meteorological conditions than reduced coal production. This update now presents a comparison of Hunter Valley coal production data and PM₁₀ concentrations measured by the UHAQMN including data for CY2023. Table 3.4 presents the Hunter Valley raw coal production between CY2013 and CY2023.

Table 3.4: Hunter Valley raw coal production (2013-2023)

Year	Hunter Valley Raw Coal Production (Mt)	Percentage of 2013 Raw Coal Production (%)
2013	158	100%
2014	160	102%
2015	146	92%
2016	145	92%
2017	146	92%
2018	151	96%
2019	155	99%
2020	146	92%
2021	142	90%
2022	126	80%
2023	136	86%

CY2013 has been nominated as a reference year to show the relative scale of coal production since the beginning of the monitoring study period. As shown in Table 3.4, the annual Hunter Valley raw coal production rates have been generally consistent over the period CY2013-CY2021, ranging between 90% and 102% of the 2013 value. However, production levels have been decreasing year on year from CY2019 through to CY2022, before increasing again in CY2023. Annual coal production rates for CY2023 are higher than CY2022 but lower than all other previous years.

Figure 3.4 presents the comparison of Hunter Valley coal production data and UHAQMN annual average PM₁₀ concentrations.




Figure 3.4: Comparison of Hunter Valley coal production data and UHAQMN annual average PM₁₀ concentrations

As shown in these data, there is no visually apparent correlation between raw coal production and ambient PM₁₀ concentrations measured by the UHAQMN between 2013 and 2019. Figure 3.4 shows inconsistency in the scale of variability in each metric across 2013 – 2019, with minor proportional variability in coal production relative to significant proportional variability in annual average PM₁₀.

From 2019 to 2022, both coal production and annual average PM₁₀ concentrations have reduced year on year. In 2023, coal production was higher than in 2022, while PM₁₀ concentration increased across the UHAQMN. The fluctuations in annual average PM₁₀ concentrations are anticipated to be related to ambient temperatures and the amount of rainfall during each of these years, rather than changes in coal production. It is recommended that relationships between PM₁₀ concentrations, coal production and rainfall continue to be reevaluated annually.

4 Closure

As noted within the previous reports and reinforced with the inclusion of CY2023 data, the temporal trends in the UHAQMN and 'Remainder of NSW' monitoring subsets show that the changes in PM₁₀ concentrations within the Upper Hunter are generally consistent with changes experienced across the rest of NSW.

With the inclusion of CY2023 data, it continues to be seen that there is a negative correlation between rainfall and particulate matter concentrations across the UHAQMN. Given the consistency between PM₁₀ trends across NSW and the UHAQMN, this relationship also holds for NSW PM₁₀ concentrations more broadly.

The differences between PM₁₀ concentrations at Background stations and Diagnostic stations are near identical across CY2013-CY2023 (up to 1 µg/m³ variability), while the range in annual average PM₁₀ concentrations across this period is 22 µg/m³. This reinforces that changes in Upper Hunter PM₁₀ concentrations are associated with regional conditions and are indicative of a minimal change in the contribution from local emission sources inclusive of mining.

There is no visually apparent correlation between raw coal production and ambient PM₁₀ concentrations measured by the UHAQMN between CY2013 and CY2019. The supporting figure shows inconsistency in the scale of variability in each metric across this period, with minor proportional variability in coal production relative to significant proportional variability in annual average PM₁₀.

From 2019 to 2022, both coal production and annual average PM₁₀ concentrations have reduced year on year. In 2023, coal production was higher than in 2022, while PM₁₀ concentration increased across the UHAQMN. The fluctuations in annual average PM₁₀ concentrations are anticipated to be related to ambient temperatures and the amount of rainfall during each of these years, rather than the reduced coal production. It is recommended that relationships between PM₁₀ concentrations, coal production and rainfall continue be reevaluated annually.

We trust that the above provides an appropriate level of detail to meet your requirements. Do not hesitate to contact the undersigned if you have any queries on the above.

Yours sincerely



Damon Roddis
Principal - Air Quality and Carbon
Zephyr Environmental
damon.roddis@zephyrenviro.com



Russ Francis
Senior Consultant – Air Quality and Carbon
Zephyr Environmental
russ.francis@zephyrenviro.com

References

BoM 2024, Bureau of Meteorology, Climate change – trends and extremes, Australian climate variability & change – Time series graphs, http://www.bom.gov.au/cgi-bin/climate/change/timeseries.cgi?graph=rain&area=nsw&season=0112&ave_yr=0, accessed February 2024

DPE 2022, Upper Hunter Air Quality Monitoring Network 2021: 5-year review 2022, <https://www.environment.nsw.gov.au/-/media/OEH/Corporate-Site/Documents/Air/upper-hunter-air-quality-monitoring-network-5-year-review-220343.pdf>, accessed February 2024

DPE, 2024, Data download facility, <https://www.airquality.nsw.gov.au/air-quality-data-services/data-download-facility>, accessed February 2024

ERM, 2020, Upper Hunter Mining Dialogue, Air Quality Monitoring Data Analysis Project, Project No.: 0526661

Zephyr, 2021, 0012 UHMD UHAQMN Annual Data Review L1 Final Update, December 2021

Zephyr, 2022, 0130 NSWMC UHAQMP Annual Data Review Final, October 2023